

CARBIDOPA AND LEVODOPA- carbidopa and levodopa tablet

Cardinal Health

CARBIDOPA and LEVODOPA TABLETS, USP

40-9182


Revised — September 2011

Rx Only

DESCRIPTION


Carbidopa and levodopa is a combination product for the treatment of Parkinson's disease and syndrome.

Carbidopa, an inhibitor of aromatic amino acid decarboxylation, is a white, crystalline compound, slightly soluble in water, with a molecular weight of 244.25. It is designated chemically as (-)-L- α -hydrazino- α -methyl- β -(3,4-dihydroxybenzene) propanoic acid monohydrate. Its molecular formula is $C_{10}H_{14}N_2O_4 \cdot H_2O$ and its structural formula is:



Tablet content is expressed in terms of anhydrous carbidopa which has a molecular weight of 226.23.

Levodopa, an aromatic amino acid, is a white, crystalline compound, slightly soluble in water, with a molecular weight of 197.19. It is designated chemically as (-)-L- α -amino- β -(3,4-dihydroxybenzene) propanoic acid. Its molecular formula is $C_9H_{11}NO_4$, and its structural formula is:



Carbidopa and levodopa tablets, for oral administration, are supplied in three strengths:

10 mg/100 mg, containing 10 mg of carbidopa and 100 mg of levodopa.

25 mg/100 mg, containing 25 mg of carbidopa and 100 mg of levodopa.

25 mg/250 mg, containing 25 mg of carbidopa and 250 mg of levodopa.

In addition, each tablet contains the following inactive ingredients:

10 mg/100 mg — Corn starch, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

25 mg/100 mg — Corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake (sunset yellow lake), magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

25 mg/250 mg — Corn starch, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

CLINICAL PHARMACOLOGY

Parkinson's disease is a progressive, neurodegenerative disorder of the extrapyramidal nervous system affecting the mobility and control of the skeletal muscular system. Its characteristic features include resting tremor, rigidity, and bradykinetic movements. Symptomatic treatments, such as levodopa therapies, may permit the patient better mobility.

Mechanism of Action

Current evidence indicates that symptoms of Parkinson's disease are related to depletion of dopamine in the corpus striatum. Administration of dopamine is ineffective in the treatment of Parkinson's disease apparently because it does not cross the blood-brain barrier. However, levodopa, the metabolic precursor of dopamine, does cross the blood-brain barrier, and presumably is converted to dopamine in the brain. This is thought to be the mechanism whereby levodopa relieves symptoms of Parkinson's disease.

Pharmacodynamics

When levodopa is administered orally it is rapidly decarboxylated to dopamine in extracerebral tissues so that only a small portion of a given dose is transported unchanged to the central nervous system. For this reason, large doses of levodopa are required for adequate therapeutic effect and these may often be accompanied by nausea and other adverse reactions, some of which are attributable to dopamine formed in extracerebral tissues.

Since levodopa competes with certain amino acids for transport across the gut wall, the absorption of levodopa may be impaired in some patients on a high protein diet.

Carbidopa inhibits decarboxylation of peripheral levodopa. It does not cross the blood-brain barrier and does not affect the metabolism of levodopa within the central nervous system.

The incidence of levodopa-induced nausea and vomiting is less with the combination product than with levodopa. In many patients, this reduction in nausea and vomiting will permit more rapid dosage titration.

Since its decarboxylase inhibiting activity is limited to extracerebral tissues, administration of carbidopa with levodopa makes more levodopa available for transport to the brain.

Pharmacokinetics

Carbidopa reduces the amount of levodopa required to produce a given response by about 75% and, when administered with levodopa, increases both plasma levels and the plasma half-life of levodopa, and decreases plasma and urinary dopamine and homovanillic acid.

The plasma half-life of levodopa is about 50 minutes, without carbidopa. When carbidopa and levodopa are administered together, the half-life of levodopa is increased to about 1.5 hours. At steady state, the bioavailability of carbidopa from carbidopa and levodopa tablets is approximately 99% relative to the concomitant administration of carbidopa and levodopa.

In clinical pharmacologic studies, simultaneous administration of carbidopa and levodopa produced greater urinary excretion of levodopa in proportion to the excretion of dopamine than administration of the two drugs at separate times.

Pyridoxine hydrochloride (vitamin B₆), in oral doses of 10 mg to 25 mg, may reverse the effects of levodopa by increasing the rate of aromatic amino acid decarboxylation. Carbidopa inhibits this action of pyridoxine; therefore, carbidopa and levodopa can be given to patients receiving supplemental pyridoxine (vitamin B₆).

INDICATIONS AND USAGE

Carbidopa and levodopa tablets are indicated in the treatment of the symptoms of idiopathic Parkinson's disease (paralysis agitans), post-encephalitic parkinsonism, and symptomatic parkinsonism which may follow injury to the nervous system by carbon monoxide intoxication and/or manganese intoxication. This product is indicated in these conditions to permit the administration of lower doses of levodopa with reduced nausea and vomiting, with more rapid dosage titration, with a somewhat smoother response, and with supplemental pyridoxine (vitamin B₆).

In some patients a somewhat smoother antiparkinsonian effect results from therapy with carbidopa and levodopa than with levodopa. However, patients with markedly irregular ("on-off") responses to levodopa have not been shown to benefit from carbidopa and levodopa.

Although the administration of carbidopa permits control of parkinsonism and Parkinson's disease with much lower doses of levodopa, there is no conclusive evidence at present that this is beneficial other than in reducing nausea and vomiting, permitting more rapid titration, and providing a somewhat smoother response to levodopa.

Certain patients who responded poorly to levodopa have improved when carbidopa and levodopa was substituted. This is most likely due to decreased peripheral decarboxylation of levodopa which results from administration of carbidopa rather than to a primary effect of carbidopa on the nervous system. Carbidopa has not been shown to enhance the intrinsic efficacy of levodopa in parkinsonian syndromes.

In considering whether to give this combination product to patients already on levodopa who have nausea and/or vomiting, the practitioner should be aware that, while many patients may be expected to improve, some do not. Since one cannot predict which patients are likely to improve, this can only be determined by a trial of therapy. It should be further noted that in controlled trials comparing carbidopa and levodopa with levodopa, about half of the patients with nausea and/or vomiting on levodopa improved spontaneously despite being retained on the same dose of levodopa during the controlled portion of the trial.

CONTRAINDICATIONS

Nonselective monoamine oxidase (MAO) inhibitors are contraindicated for use with carbidopa and levodopa. These inhibitors must be discontinued at least two weeks prior to initiating therapy with this combination product. Carbidopa and levodopa may be administered concomitantly with the manufacturer's recommended dose of an MAO inhibitor with selectivity for MAO type B (e.g., selegiline HCl) (See PRECAUTIONS, Drug Interactions).

Carbidopa and levodopa is contraindicated in patients with known hypersensitivity to any component of this drug, and in patients with narrow-angle glaucoma.

Because levodopa may activate a malignant melanoma, carbidopa and levodopa should not be used in patients with suspicious, undiagnosed skin lesions or a history of melanoma.

WARNINGS

When carbidopa and levodopa tablets are to be given to patients who are being treated with levodopa, levodopa must be discontinued at least twelve hours before therapy with the combination product is started. In order to reduce adverse reactions, it is necessary to individualize therapy. See DOSAGE AND ADMINISTRATION section before initiating therapy.

The addition of carbidopa with levodopa in the form of this combination product reduces the peripheral effects (nausea, vomiting) due to decarboxylation of levodopa; however, carbidopa does not decrease the adverse reactions due to the central effects of levodopa. Because carbidopa permits more levodopa to reach the brain and more dopamine to be formed, certain adverse CNS effects, e.g., dyskinesias (involuntary movements), may occur at lower dosages and sooner with carbidopa and levodopa than with levodopa alone.

Levodopa alone, as well as carbidopa and levodopa, is associated with dyskinesias. The occurrence of dyskinesias may require dosage reduction.

As with levodopa, the combination product may cause mental disturbances. These reactions are thought to be due to increased brain dopamine following administration of levodopa. All patients should be observed carefully for the development of depression with concomitant suicidal tendencies. Patients with past or current psychoses should be treated with caution.

Carbidopa and levodopa should be administered cautiously to patients with severe cardiovascular or pulmonary disease, bronchial asthma, renal, hepatic or endocrine disease.

As with levodopa, care should be exercised in administering the combination product, to patients with a history of myocardial infarction who have residual atrial, nodal, or ventricular arrhythmias. In such patients, cardiac function should be monitored with particular care during the period of initial dosage adjustment, in a facility with provisions for intensive cardiac care.

As with levodopa, treatment with the combination product may increase the possibility of upper gastrointestinal hemorrhage in patients with a history of peptic ulcer.

Neuroleptic Malignant Syndrome (NMS):

Sporadic cases of a symptom complex resembling NMS have been reported in association with dose reductions or withdrawal of therapy with carbidopa and levodopa. Therefore, patients should be observed carefully when the dosage of carbidopa and levodopa is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia. Neurological findings, including muscle rigidity, involuntary movements, altered consciousness, mental status changes; other disturbances, such as autonomic dysfunction, tachycardia, tachypnea, sweating, hyper- or hypotension; laboratory findings, such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin have been reported.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complex if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) intensive symptomatic treatment and medical monitoring and 2) treatment of any concomitant serious medical problems for which specific treatments are available. Dopamine agonists, such as bromocriptine, and muscle relaxants, such as dantrolene, are often used in the treatment of NMS, however, their effectiveness has not been demonstrated in controlled studies.

PRECAUTIONS

General

As with levodopa, periodic evaluations of hepatic, hematopoietic, cardiovascular, and renal function are recommended during extended therapy.

Patients with chronic wide-angle glaucoma may be treated cautiously with carbidopa and levodopa provided the intraocular pressure is well-controlled and the patient is monitored carefully for changes in intraocular pressure during therapy.

Dopaminergic agents, including levodopa, may be associated with somnolence and very rarely episodes of sudden onset of sleep. In some cases, these episodes may occur without awareness or warning during daily activities. Patients must be informed of this and advised to exercise caution while driving or operating machines while being treated with dopaminergic agents, including levodopa. Patients who

have experienced somnolence and/or an episode of sudden sleep onset must refrain from driving or operating machines (see Information for Patients).

Melanoma

Epidemiological studies have shown that patients with Parkinson's disease have a higher risk (2- to approximately 6-fold higher) of developing melanoma than the general population. Whether the increased risk observed was due to Parkinson's disease or other factors, such as drugs used to treat Parkinson's disease, is unclear.

For the reasons stated above, patients and providers are advised to monitor for melanomas frequently and on a regular basis when using carbidopa and levodopa for any indication. Ideally, periodic skin examinations should be performed by appropriately qualified individuals (e.g., dermatologists)

Information for Patients

The patient should be informed that this combination product is an immediate-release formulation of carbidopa and levodopa that is designed to begin release of ingredients within 30 minutes. It is important that carbidopa and levodopa be taken at regular intervals according to the schedule outlined by the physician. The patient should be cautioned not to change the prescribed dosage regimen and not to add any additional antiparkinson medications, including other carbidopa and levodopa preparations, without first consulting the physician.

Patients should be advised that sometimes a 'wearing-off' effect may occur at the end of the dosing interval. The physician should be notified if such response poses a problem to life style.

Patients should be advised that occasionally, dark color (red, brown, or black) may appear in saliva, urine, or sweat after ingestion of carbidopa and levodopa. Although the color appears to be clinically insignificant, garments may become discolored.

The patient should be advised that a change in diet to foods that are high in protein may delay the absorption of levodopa and may reduce the amount taken up in the circulation. Excessive acidity also delays stomach emptying, thus delaying the absorption of levodopa. Iron salts (such as in multi vitamin tablets) may also reduce the amount of levodopa available to the body. The above factors may reduce the clinical effectiveness of the levodopa or carbidopa and levodopa therapy.

Patients should be alerted to the possibility of sudden onset of sleep during daily activities, in some cases without awareness or warning signs, when they are taking dopaminergic agents, including levodopa. Patients should be advised to exercise caution while driving or operating machinery and that if they have experienced somnolence and/or sudden sleep onset, they must refrain from these activities. (See PRECAUTIONS: General.)

There have been reports of patients experiencing intense urges to gamble, increased sexual urges, and other intense urges, and the inability to control these urges while taking one or more of the medications that increase central dopaminergic tone and that are generally used for the treatment of Parkinson's disease, including carbidopa and levodopa. Although it is not proven that the medications caused these events, these urges were reported to have stopped in some cases when the dose was reduced or the medication was stopped. Prescribers should ask patients about the development of new or increased gambling urges, sexual urges or other urges while being treated with carbidopa and levodopa. Patients should inform their physician if they experience new or increased gambling urges, increased sexual urges, or other intense urges while taking carbidopa and levodopa. Physicians should consider dose reduction or stopping the medication if a patient develops such urges while taking carbidopa and levodopa.

Laboratory Tests

Abnormalities in laboratory tests may include elevations of liver function tests such as alkaline phosphatase, SGOT (AST), SGPT (ALT), lactic dehydrogenase, and bilirubin. Abnormalities in blood urea nitrogen and positive Coombs test have also been reported. Commonly, levels of blood urea

nitrogen, creatinine, and uric acid are lower during administration of this combination product than with levodopa.

Carbidopa and levodopa may cause a false-positive reaction for urinary ketone bodies when a test tape is used for determination of ketonuria. This reaction will not be altered by boiling the urine specimen. False-negative tests may result with the use of glucose-oxidase methods of testing for glucosuria.

Cases of falsely diagnosed pheochromocytoma in patients on carbidopa and levodopa therapy have been reported very rarely. Caution should be exercised when interpreting the plasma and urine levels of catecholamines and their metabolites in patients on levodopa or carbidopa and levodopa therapy.

Drug Interactions

Caution should be exercised when the following drugs are administered concomitantly with carbidopa and levodopa.

Symptomatic postural hypotension has occurred when carbidopa and levodopa was added to the treatment of a patient receiving antihypertensive drugs. Therefore, when therapy with carbidopa and levodopa is started, dosage adjustment of the antihypertensive drug may be required.

For patients receiving MAO inhibitors (Type A or B), see CONTRAINDICATIONS. Concomitant therapy with selegiline and carbidopa and levodopa may be associated with severe orthostatic hypotension not attributable to carbidopa and levodopa alone (see CONTRAINDICATIONS).

There have been rare reports of adverse reactions, including hypertension and dyskinesia, resulting from the concomitant use of tricyclic antidepressants and carbidopa and levodopa.

Dopamine D₂ receptor antagonists (e.g., phenothiazines, butyrophenones, risperidone) and isoniazid may reduce the therapeutic effects of levodopa. In addition, the beneficial effects of levodopa in Parkinson's disease have been reported to be reversed by phenytoin and papaverine. Patients taking these drugs with carbidopa and levodopa should be carefully observed for loss of therapeutic response.

Iron salts may reduce the bioavailability of levodopa and carbidopa. The clinical relevance is unclear.

Although metoclopramide may increase the bioavailability of levodopa by increasing gastric emptying, metoclopramide may also adversely affect disease control by its dopamine receptor antagonistic properties.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a two-year bioassay of carbidopa and levodopa, no evidence of carcinogenicity was found in rats receiving doses of approximately two times the maximum daily human dose of carbidopa and four times the maximum daily human dose of levodopa.

In reproduction studies with carbidopa and levodopa, no effects on fertility were found in rats receiving doses of approximately two times the maximum daily human dose of carbidopa and four times the maximum daily human dose of levodopa.

Pregnancy

Pregnancy Category C: No teratogenic effects were observed in a study in mice receiving up to 20 times the maximum recommended human dose of carbidopa and levodopa. There was a decrease in the number of live pups delivered by rats receiving approximately two times the maximum recommended human dose of carbidopa and approximately five times the maximum recommended human dose of levodopa during organogenesis. Carbidopa and levodopa caused both visceral and skeletal malformations in rabbits at all doses and ratios of carbidopa/levodopa tested, which ranged from 10 times/5 times the maximum recommended human dose of carbidopa/levodopa to 20 times/10 times the maximum recommended human dose of carbidopa/levodopa.

There are no adequate or well-controlled studies in pregnant women. It has been reported from individual cases that levodopa crosses the human placental barrier, enters the fetus, and is metabolized.

Carbidopa concentrations in fetal tissue appeared to be minimal. Use of carbidopa and levodopa in women of childbearing potential requires that the anticipated benefits of the drug be weighed against possible hazards to mother and child.

Nursing Mothers

In a study of one nursing mother with Parkinson's disease, excretion of levodopa in human breast milk was reported. Therefore, caution should be exercised when carbidopa and levodopa is administered to a nursing woman.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established. Use of the drug in patients below the age of 18 is not recommended.

ADVERSE REACTIONS

The most common adverse reactions reported with carbidopa and levodopa have included dyskinesias, such as choreiform, dystonic, and other involuntary movements and nausea.

The following other adverse reactions have been reported with carbidopa and levodopa:

Body As A Whole: chest pain, asthenia.

Cardiovascular: cardiac irregularities, hypotension, orthostatic effects including orthostatic hypotension, hypertension, syncope, phlebitis, palpitation.

Gastrointestinal: dark saliva, gastrointestinal bleeding, development of duodenal ulcer, anorexia, vomiting, diarrhea, constipation, dyspepsia, dry mouth, taste alterations.

Hematologic: agranulocytosis, hemolytic and non-hemolytic anemia, thrombocytopenia, leukopenia.

Hypersensitivity: angioedema, urticaria, pruritus, Henoch-Schonlein purpura, bullous lesions (including pemphigus-like reactions).

Musculoskeletal: back pain, shoulder pain, muscle cramps.

Nervous System/Psychiatric: psychotic episodes including delusions, hallucinations, and paranoid ideation, neuroleptic malignant syndrome (see WARNINGS), bradykinetic episodes ("on-off" phenomenon), confusion, agitation, dizziness, somnolence, dream abnormalities including nightmares, insomnia, paresthesia, headache, depression with or without development of suicidal tendencies, dementia, Pathological gambling, increased libido including hypersexuality, impulse control symptoms. Convulsions also have occurred; however, a causal relationship with carbidopa and levodopa has not been established.

Respiratory: dyspnea, upper respiratory infection.

Skin: rash, increased sweating, alopecia, dark sweat.

Urogenital: urinary tract infection, urinary frequency, dark urine.

Laboratory Tests

decreased hemoglobin and hematocrit; abnormalities in alkaline phosphatase, SGOT (AST), SGPT (ALT), lactic dehydrogenase, bilirubin, blood urea nitrogen (BUN), Coombs test; elevated serum glucose; white blood cells, bacteria, and blood in the urine.

Other adverse reactions that have been reported with levodopa alone and with various carbidopa and levodopa formulations, and may occur with this combination product are:

Body As A Whole: abdominal pain and distress, fatigue.

Cardiovascular: myocardial infarction.

Gastrointestinal: gastrointestinal pain, dysphagia, sialorrhea, flatulence, bruxism, burning sensation of the tongue, heartburn, hiccups.

Metabolic: edema, weight gain, weight loss.

Musculoskeletal: leg pain.

Nervous System/Psychiatric: ataxia, extrapyramidal disorder, falling, anxiety, gait abnormalities, nervousness, decreased mental acuity, memory impairment, disorientation, euphoria, blepharospasm (which may be taken as an early sign of excess dosage; consideration of dosage reduction may be made at this time), trismus, increased tremor, numbness, muscle twitching, activation of latent Horner's syndrome, peripheral neuropathy.

Respiratory: pharyngeal pain, cough.

Skin: malignant melanoma (see also CONTRAINDICATIONS), flushing.

Special Senses: oculogyric crises, diplopia, blurred vision, dilated pupils.

Urogenital: urinary retention, urinary incontinence, priapism.

Miscellaneous: bizarre breathing patterns, faintness, hoarseness, malaise, hot flashes, sense of stimulation.

Laboratory Tests: decreased white blood cell count and serum potassium; increased serum creatinine and uric acid; protein and glucose in urine.

OVERDOSAGE

Management of acute overdosage with carbidopa and levodopa is the same as management of acute overdosage with levodopa. Pyridoxine is not effective in reversing the actions of this product.

General supportive measures should be employed, along with immediate gastric lavage. Intravenous fluids should be administered judiciously and an adequate airway maintained. Electrocardiographic monitoring should be instituted and the patient carefully observed for the development of arrhythmias; if required, appropriate antiarrhythmic therapy should be given. The possibility that the patient may have taken other drugs as well as carbidopa and levodopa tablets should be taken into consideration. To date, no experience has been reported with dialysis; hence, its value in overdosage is not known.

Based on studies in which high doses of levodopa and/or carbidopa were administered, a significant proportion of rats and mice given single oral doses of levodopa of approximately 1500-2000 mg/kg are expected to die. A significant proportion of infant rats of both sexes are expected to die at a dose of 800 mg/kg. A significant proportion of rats are expected to die after treatment with similar doses of carbidopa. The addition of carbidopa in a 1:10 ratio with levodopa increases the dose at which a significant proportion of mice are expected to die to 3360 mg/kg.

DOSAGE AND ADMINISTRATION

The optimum daily dosage of carbidopa and levodopa must be determined by careful titration in each patient. Carbidopa and levodopa tablets are available in a 1:4 ratio of carbidopa to levodopa (25 mg/100 mg) as well as 1:10 ratio (25 mg/250 mg and 10 mg/100 mg). Tablets of the two ratios may be given separately or combined as needed to provide the optimum dosage.

Studies show that peripheral dopa decarboxylase is saturated by carbidopa at approximately 70 to 100 mg a day. Patients receiving less than this amount of carbidopa are more likely to experience nausea and vomiting.

Usual Initial Dosage: Dosage is best initiated with one tablet of carbidopa and levodopa 25 mg/100 mg three times a day. This dosage schedule provides 75 mg of carbidopa per day. Dosage

may be increased by one tablet every day or every other day, as necessary, until a dosage of eight tablets of carbidopa and levodopa 25 mg/100 mg a day is reached.

If carbidopa and levodopa 10 mg/100 mg is used, dosage may be initiated with one tablet three or four times a day. However, this will not provide an adequate amount of carbidopa for many patients. Dosage may be increased by one tablet every day or every other day until a total of eight tablets (2 tablets q.i.d.) is reached.

How To Transfer Patients From Levodopa: Levodopa must be discontinued at least twelve hours before starting this combination product. A daily dosage of carbidopa and levodopa should be chosen that will provide approximately 25% of the previous levodopa dosage. Patients who are taking less than 1500 mg of levodopa a day should be started on one tablet of carbidopa and levodopa 25 mg/100 mg three or four times a day. The suggested starting dosage for most patients taking more than 1500 mg of levodopa is one tablet of carbidopa and levodopa 25 mg/250 mg three or four times a day.

Maintenance: Therapy should be individualized and adjusted according to the desired therapeutic response. At least 70 to 100 mg of carbidopa per day should be provided. When a greater proportion of carbidopa is required, one 25 mg/100 mg tablet may be substituted for each 10 mg/100 mg tablet. When more levodopa is required, each 25 mg/250 mg tablet should be substituted for a 25 mg/100 mg tablet or a 10 mg/100 mg tablet. If necessary, the dosage of carbidopa and levodopa 25 mg/250 mg may be increased by one-half or one tablet every day or every other day to a maximum of eight tablets a day. Experience with total daily dosages of carbidopa greater than 200 mg is limited.

Because both therapeutic and adverse responses occur more rapidly with this combination product than with levodopa alone, patients should be monitored closely during the dose adjustment period. Specifically, involuntary movements will occur more rapidly with carbidopa and levodopa than with levodopa. The occurrence of involuntary movements may require dosage reduction. Blepharospasm may be a useful early sign of excess dosage in some patients.


Addition Of Other Antiparkinsonian Medications: Standard drugs for Parkinson's disease, other than levodopa without a decarboxylase inhibitor, may be used concomitantly while carbidopa and levodopa is being administered, although dosage adjustments may be required.


Interruption Of Therapy: Sporadic cases of a symptom complex resembling Neuroleptic Malignant Syndrome (NMS) have been associated with dose reductions and withdrawal of carbidopa and levodopa. Patients should be observed carefully if abrupt reduction or discontinuation of carbidopa and levodopa is required, especially if the patient is receiving neuroleptics. (See WARNINGS.)


If general anesthesia is required, carbidopa and levodopa may be continued as long as the patient is permitted to take fluids and medication by mouth. If therapy is interrupted temporarily, the patient should be observed for symptoms resembling NMS, and the usual daily dosage may be administered as soon as the patient is able to take oral medication.

HOW SUPPLIED

Carbidopa and Levodopa Tablets USP are supplied as follows:

10 mg/100 mg — Each dark blue, mottled, round tablet imprinted with  on one side and 538 and bisect on the other contains 10 mg of Carbidopa and 100 mg of Levodopa and is supplied in cards of 30 tablets (NDC 55154-3829-9).

25 mg/100 mg — Each yellow, mottled, round tablet imprinted with  on one side and 539 and bisect on the other contains 25 mg of Carbidopa and 100 mg of Levodopa and is supplied in cards of 28 tablets (NDC 55154-3828-3) and in cards of 30 tablets (NDC 55154-3828-9).

25 mg/250 mg — Each light blue, mottled, round tablet imprinted with  on one side and 540 and bisect on the other contains 25 mg of Carbidopa and 250 mg of Levodopa and is supplied in cards of 30 tablets (NDC 55154-3830-9).

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Protect from light.

Dispense in a well-closed, light-resistant container as defined in the USP.

40-9182

Revised — September 2011

Repackaged by

Cardinal Health

Zanesville, OH 43701

IT47854160415

Package/Label Display Panel

Carbidopa and Levodopa

10 mg/ 100 mg

tablets, USP

CARBIDOPA AND LEVODOPA
10 mg/100 mg
tablets, USP

LOT: xxxxxxxxxx EXP: xx/xx/xx

LTC49413810214



CUSTOMER ID

Package/Label Display Panel

Carbidopa and Levodopa

10 mg/ 100 mg tablets, USP

QTY 30

NDC 55154-3829-9 (CARDINAL HEALTH)
CARBIDOPA AND LEVODOPA
10 mg/100 mg tablets, USP
QTY 30 RX ONLY

Each Tablet Contains: Carbidopa, USP 10 mg*
 *(Anhydrous equivalent)
 Levodopa, USP 100 mg

Usual Adult Dosage: See product insert for prescribing
 information, precautions and warnings.

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).
 [See USP Controlled Room Temperature].





























Protect from light.

Dispense in a well-closed, light-resistant container as defined in the USP.

This package is intended for in-patient institutional use only.
 Keep this and all drugs out of the reach of children.

Manufactured by:
 Actavis Elizabeth LLC
 Elizabeth, NJ 07207 USA
 Distributed by: Actavis Pharma, Inc.
 Parsippany, NJ 07054 USA

Repackaged By: Cardinal Health
 Zanesville, OH 43701
 LTB49413810415

Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 10 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3829-9 RPKD BY: Cardinal Health Zanesville OH 43701

Package/Label Display Panel

Carbidopa and Levodopa

25 mg/ 100 mg

tablets, USP

CARBIDOPA AND LEVODOPA
25 mg/100 mg
tablets, USP

LOT: xxxxxxxxxx EXP: xx/xx/xx

LTC47857210814



CUSTOMER ID

Package/Label Display Panel

Carbidopa and Levodopa

25 mg/ 100 mg

tablets, USP

QTY 28

NDC 55154-3828-3 (CARDINAL HEALTH)
CARBIDOPA AND LEVODOPA
25 mg/100 mg
tablets, USP
QTY 28 RX ONLY

Each Tablet Contains: Carbidopa, USP 25 mg*
 *(Anhydrous equivalent)
 Levodopa, USP 100 mg

Contains FD&C Yellow #6 Aluminum Lake
 (Sunset Yellow Lake) as a color additive.

Usual Adult Dosage: See product insert for
 prescribing information, precautions and warnings.


Store at 25°C (77°F); excursions
 permitted to 15° to 30°C (59° to 86°F).
 [See USP Controlled Room Temperature].
 Protect from light.
 Dispense in a well-closed, light-resistant
 container as defined in the USP.

This package is intended for in-patient institutional use only.
 Keep this and all drugs out of the reach of children.

Manufactured by: Actavis Elizabeth LLC
 Elizabeth, NJ 07207 USA

Distributed by: Actavis Pharma, Inc.
 Parsippany, NJ 07054 USA

Repackaged By: Cardinal Health
 Zanesville, OH 43701
 LTB49457910715

Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701
Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701	Carbidopa and Levodopa tablet, USP 25 mg/100 mg LOT: XXXXXXXXXX EXP: XX/XX/XX  55154-3828-3 RPKD BY: Cardinal Health Zanesville OH 43701

PRINCIPAL DISPLAY PANEL

Carbidopa and Levodopa

25 mg/ 100 mg

tablets, USP

**CARBIDOPA AND LEVODOPA
25 mg/100 mg
tablets, USP**

LOT: xxxxxxxxxx EXP: xx/xx/xx

LTC47857210814



CUSTOMER ID

PRINCIPAL DISPLAY PANEL

Carbidopa and Levodopa

25 mg/ 100 mg tablets, USP

Qty 30

RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa

25 mg/ 250 mg

tablets, USP

CARBIDOPA AND LEVODOPA

25 mg/250 mg

tablets, USP

LOT: xxxxxxxxxx EXP: xx/xx/xx

LTC49442290214



CUSTOMER ID


Package/Label Display Panel

Carbidopa and Levodopa

25 mg/ 250 mg tablets, USP

QTY 30

NDC 55154-3830-9 (CARDINAL HEALTH)
CARBIDOPA AND LEVODOPA
25 mg/250 mg tablets, USP
QTY 30 RX ONLY



Each Tablet Contains: Carbidopa, USP 25 mg*
*(Anhydrous equivalent)
Levodopa, USP 250 mg

Usual Adult Dosage: See product insert for
prescribing information, precautions and warnings.
Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).
[See USP Controlled Room Temperature].

Protect from light.

Dispense in a well-closed, light-resistant
container as defined in the USP.

This package is intended for in-patient institutional use only.
Keep this and all drugs out of the reach of children.

Manufactured by:
Actavis Elizabeth LLC
Elizabeth, NJ 07207 USA
Distributed by: Actavis Pharma, Inc.
Parsippany, NJ 07054 USA

Repackaged By: Cardinal Health
Zanesville, OH 43701
LTB49442290415

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

Carbidopa and Levodopa
tablet, USP
25 mg/250 mg
LOT: XXXXXXXXXXXX EXP: XX/XX/XX
55154-3830-9
RPKD BY: Cardinal Health
Zanesville OH 43701

carbidopa and levodopa tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55154-3829(NDC:0228-2538)
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CARBIDOPA (UNII: MNX7R8C5VO) (CARBIDOPA ANHYDROUS - UNII:KR87B45RGH)	CARBIDOPA ANHYDROUS	10 mg
LEVODOPA (UNII: 46627O600J) (LEVODOPA - UNII:46627O600J)	LEVODOPA	100 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

Product Characteristics			
Color	BLUE (dark blue, mottled)	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	R;538
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-3829-9	30 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074260	09/01/1993	

CARBIDOPA AND LEVODOPA

carbidopa and levodopa tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55154-3828(NDC:0228-2539)

Route of Administration		ORAL	DEA Schedule	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
CARBIDOPA (UNII: MNX7R8C5VO) (CARBIDOPA ANHYDROUS - UNII:KR87B45RGH)			CARBIDOPA ANHYDROUS	25 mg
LEVODOPA (UNII: 46627O600J) (LEVODOPA - UNII:46627O600J)			LEVODOPA	100 mg
Inactive Ingredients				
Ingredient Name				Strength
STARCH, CORN (UNII: O8232NY3SJ)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
Product Characteristics				
Color	YELLOW (mottled)		Score	2 pieces
Shape	ROUND		Size	9mm
Flavor			Imprint Code	R;539
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-3828-9	30 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55154-3828-3	28 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA074260	09/01/1993	

CARBIDOPA AND LEVODOPA

carbidopa and levodopa tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55154-3830(NDC:0228-2540)
Route of Administration	ORAL	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

CARBIDOPA (UNII: MNX7R8C5VO) (CARBIDOPA ANHYDROUS - UNII:KR87B45RGH)		CARBIDOPA ANHYDROUS	25 mg	
LEVODOPA (UNII: 46627O600J) (LEVODOPA - UNII:46627O600J)		LEVODOPA	250 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
Product Characteristics				
Color	BLUE (light blue, mottled)	Score	2 pieces	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	R;540	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-3830-9	30 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA074260		09/01/1993	

Labeler - Cardinal Health (188557102)

Establishment			
Name	Address	ID/FEI	Business Operations
Cardinal Health		188557102	REPACK(55154-3828, 55154-3830, 55154-3829)